



Effect of Methanolic Extract of *Securigera securidaca* as Antioxidant and Antibacterial Activities

Ghassab M. Al-Mazaideh^{1*} and Saleh A. Al-Quran²

¹Department of Chemistry and Chemical Technology, Faculty of Science, Tafila Technical University, P.O.Box 179, Tafila 66110, Jordan.

²Department of Biological Sciences, Faculty of Science, Mutah University, P.O.Box 7, Mutah 61710, Al-Karak, Jordan.

Authors' contributions

This work was carried out in collaboration between both authors. Author GMA designed the study, carried out all experimental work and wrote the first draft of the manuscript. Author SAA managed the analyses of the study and managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2020/v32i1130498

Editor(s):

(1) Dr. Vasudevan Mani, Qassim University, Saudi Arabia.

Reviewers:

(1) Dalia Mahmood, Al-Nahrain University, Iraq.

(2) Mukesh K. Berwal, ICAR-Central Institute for Arid Horticulture, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/58374>

Original Research Article

Received 20 April 2020

Accepted 27 June 2020

Published 06 July 2020

ABSTRACT

In the present work, the phytochemical screening, polyphenolic content, antibacterial activity and antioxidant activity of *Securigera securidaca* seeds in methanol were carried out. Phytochemical analysis of seeds showed the presence of alkaloids, flavonoids, saponins, terpenoids, steroids and glycosides. Total phenolic content was estimated by Folin Ciocalteu method and the result showed the highest phenolic content of 62.28 mg/g. Methanolic extract was screened for antibacterial activity by disc diffusion method and it found to be potent. The MIC of methanol extract identified by broth dilution method showed a MIC value of 0.25 mg/ml for both *E. coli* and *Kl. Oxytoca*, and also 0.5 mg/ml for both *S. aureus* and *S. epidermis*. The antioxidant effect of the seeds was tested by DPPH scavenging activity as *in vitro* assay. The extract had potent inhibitory activity (IC₅₀) value of 0.057 mg/ml. The finding experimental results showed that methanolic extract of *Securigera securidaca* is important as a source of antibacterial activity and polyphenolic antioxidants.

*Corresponding author: E-mail: gmazideh@ttu.edu.jo, gsuadi2016@gmail.com;

Keywords: *Securigera securidaca*; antioxidant; anticorrosion; polyphenolic content; MIC.

1. INTRODUCTION

Medicinal plants have been known as green anticorrosion inhibitors [1-6], analgesics, antimicrobial, antioxidant, antispasmodics, and diuretics [7] since ancient times. Therefore, it becomes important starting material of drugs due to their content of secondary metabolites in plant. Currently, new antibacterial and antioxidant agents that have lesser side effects and better efficacy is promising approach in order to treat infectious diseases.

Securigera securidaca (*S. securidaca*) is an annual herb occurring wild in West Asia, Africa and Europe. Also called *goat pea* and popularly names as *Gandeh Talkheh* [8]. Several experimental studies have shown beneficial effects of *S. securidaca* seeds as natural promise agents for epilepsy in Iranian folk medicine [9], enhancing antidiabetic, chronotropic, treatment of disorders such as hyperlipidemia, hypoglycemic effects, diuretic, hypokalaemic activities [4] and as anti-HIV-1 activity [10]. Moreover, experimental studies showed its role in reducing the level of cholesterol and triglyceride in serum of high-fat fed rats [11,12]. It is reported that the ethanolic and aqueous extract of *S. securidaca* contains various classes of secondary metabolites such as steroids,

flavonoids, alkaloids, tannins, cardenolides and penta cyclic triterpenoid type saponins [13,14]. Since these compounds may have the potential to inhibit Gram positive and negative bacteria and act as potent antioxidants, they have the ideal chemical structure for scavenging free radicals [15]. The petroleum ether extract of *S. securidaca* seeds showed antibacterial activities against *Staphylococcus aureus* (*S. aureus*) and *Pseudomonas aeruginosa* (*P. aeruginosa*) while the chloroform extract showed inhibitory effect only for *S. aureus*. The odd thing is, no antimicrobial activity of methanol extract, although there are many important natural compounds in the crude of *S. securidaca* seeds and there are many amino acids that have been determined in the composition of *Securigera securidaca* seeds such as Alanine, asparagine, arginine, aspartic acid, glutamic acid, valine and tyrosine [16,17].

Thus, this study (Fig. 1) aims to evaluate the effect of methanolic extract of *S. securidaca* seeds which will be published for the first time as antibacterial activity against *Escherichia coli* (*E. coli*), *S. aureus* (*S. aureus*), *Staphylococcus epidermis* (*S. epidermis*) and *Klebsiella oxytoca* (*Kl. oxytoca*), also antioxidant effects in order to use it in some infectious diseases.




Fig. 1. The overall work on methanolic extract of *S. securidaca* seeds

2. MATERIAL AND METHODS

2.1 Collection and Preparation of the Plant Materials

S. securidaca seeds brought from local herbal shop (Karak- Jordan) in the summer of 2017.

The seeds were grounded to be fine powder using a coffee blender and stored in the special container until use.

2.2 Preparation of Methanolic Extract

Twenty-five grams of seeds powder were soaked in 250 ml of 96% methanol and it was put in the shaker device at 150 rpm, in dark place for four days at room temperature and stored in a refrigerator for three days. The extract was then filtered using a Buchner funnel under vacuum. The filtrate was centrifuged at 3000 rpm for 15 minutes, and then extract concentrated in the rotary evaporator under vacuum at 50°C. The crude was left in open vials in the fume hood for four days at room temperature and stored thereafter at 4°C in a glass container until further use [18,19].

2.3 Extraction Yield

The yield of crude methanol extract was calculated (% , W_1/W_2) as:

$$\text{Yield} = W_1/W_2 \times 100\%$$

Where W_1 is the weight of dried and ground plant material after evaporation of methanol and W_2 is the weight of powdered plant.

2.4 Qualitative Phytochemical Analysis

Phytochemical screening of primary and secondary metabolic compounds such as alkaloids, tannins, steroids, terpenoids, saponin glycosides, flavonoids, volatile oils, starch, phenols and proteins were conducted on seed extract according to standard phytochemical methods [20,21].

2.5 Determination of Total Phenolic Content (TPC) in the Methanolic Crude

The Folin-Ciocalteu assay method [22] was used to determine the total soluble phenolic content in plant extract in terms of Gallic acid. In this experiment: plant extract (0.2 ml, three replicates); 1 ml of Folin-Ciocalteu reagent was introduced into test tubes; The mixtures were

neutralized with 0.8 ml of 7.5 % of Na_2CO_3 and the final concentration of the plant extract in the solution was 500 $\mu\text{g/ml}$. The tubes were mixed and shaken well to allow for reaction and left for 30 minutes at room temperature for color development and the absorbance was measured at 760 nm using a spectrophotometer. TPC was calculated according to the standard calibration curve of Gallic acid (GA) solutions at different concentration (0 to 25 $\mu\text{g/ml}$). TPC was expressed as Gallic acid equivalents (GAE) in milligrams per gram plant extract [23].

2.6 Evaluation of Antibacterial Activity

2.6.1 Microorganism and growth conditions

The extraction was tested for antibacterial activity *in vitro* using both Gram negative and Gram positive bacteria (four microorganism including *E. coli*, *S. aureus*, *S. epidermis* and *Kl. oxytoca*). These stock cultures of bacteria were obtained from research Lab., Department of Biology, Mutah University. Antimicrobial activity evaluations were performed using the agar disc diffusion method [24,25]. TPZ (10 μg) was employed as positive control, whereas a negative control contains pure methanol or distilled water. The plates were incubated at 37°C for 18 – 24 h. The antimicrobial activity of the extract was determined by measuring the diameter of inhibition zone (mm) against each bacterium. The tests were performed in triplicate and reported as mean \pm standard deviation (SD).

2.6.2 Minimum inhibitory concentration (MIC)

Broth dilution method [26] was used to determine MIC. The extract dissolved in 10% DMSO in methanol was first diluted to the highest concentration (200 mg/ml) to be tested and then six-fold serial dilution was made in the concentration range of 0.0625 - 200 mg/ml. The extract solutions were added to a nutrient broth in separate test tubes inoculated with the respective standardized suspension of a strain adjusted to a concentration of 1×10^8 colony/ml. Each tube contains various extract at concentration of 0 (control), 0.0625, 0.125, 0.250, 0.500, 1 and 2 mg/ml in broth medium. These tubes were incubated at 37°C overnight and observed for visible growth (turbidity). MIC can be determined by examining broth tubes compared to control tube, containing only broth and inoculums without extract. Tubes that remain clear indicate no active growth and show the lowest concentration of extract, it was considered as MIC.

2.7 Evaluation of Antioxidant Activity of the Extracts

2.7.1 DPPH radical scavenging activity

The free radical scavenging activity of crude extract was estimated based on the previous reported procedure using the stable 2, 2-diphenyl-1-picrylhydrazyl radical, known as DPPH [27]. Briefly, different concentrations of the extracts were mixed with 5 ml of 0.004% methanol solution of DPPH (plant concentration in the solution varies from 0 to 2000 µg/mL). The mixture was shaken vigorously and left to stand for 30 min with incubation at 37°C. After that, the absorbance of the resulting mixture (DPPH with extract) was read against methanol at 517 nm using a spectrophotometer. All determinations were done at least in triplicate. The radical scavenging activity (capability to scavenge the DPPH radical) was calculated as a percentage of DPPH discoloration using the following equation:

$$\text{Percentage of DPPH discoloration} = [(A_c - A_s)/A_c] \times 100 (\%)$$

Where A_c is the absorbance of the control reaction (DPPH solution without the tested extract), and A_s is the sample is the absorbance of the presence of all of the extract samples and reagents. IC_{50} (crude concentration providing 50% inhibition) was determined by a graph plotting the percentage inhibition against crude concentration. Trolox equivalent per gram dry weight can be calculated by creating a standard curve of Trolox standards (concentration 0 to 1.5 µg/ml) versus their absorbance. This curve was used to be a standard for the construction of the calibration curve, and the percentage of DPPH discoloration was expressed as mg Trolox equivalents per gram of plant extract [28].

2.8 Statistical Analysis

All Experimental data were recorded in triplicate and the results were expressed as a mean \pm standard deviation (SD). The IC_{50} value was calculated by Microsoft Excel 2010.

3. RESULTS

3.1 Phytochemical Screening

Qualitative phytochemical analysis of the phytochemical constituents presents in methanolic extract of *S. securidaca* seeds

showed the presence of alkaloids, flavonoids and saponins in high content seeds extract; terpenoids, steroids and glycosides in mild content of the extract (Table 1).

Table 1. Phytochemical analysis for the methanolic extract of *S. securidaca* seeds

Phytochemical constituents	Methanolic extract
Alkaloids	+++
Flavonoids	+++
Terpenoids	++
Steroids	++
Glycosides	++
Saponins	+++

Where: (++) means mild content, and (+++) high content

3.2 Total Phenolic Content

Folin Ciocalteu's method and the Gallic acid (GA) were used as standard compounds to find the total phenolic content (TPC). The TPC value of methanolic extract of *S. securidaca* seed was calculated using the standard curve equation of Gallic acid equivalent (GAE) in mg/g plant extract.

$$Y = 0.0469X + 0.2703, R^2 = 0.9601 \quad \text{Eq. (1)}$$

Where Y is the absorbance at 760 nm and X is the amount of total phenolics in the plant extract. The TPC of the plant crude was 62.28 mg GAE/g of plant extract.

3.3 Antimicrobial Effects

Table 2 shows the antibacterial activity of methanolic extract of the plant against four bacterial strains including *E. coli*, *S. aureus*, *S. epidermis* and *Kl. Oxytoca*. The extraction has an inhibitory effect on the growth of bacterial strains in disc diffusion method and in agar well diffusion method with an inhibition zone from 4 mm to 13 mm diameters at different extract concentration (Fig. 2).

Table 3 summarizes the MIC results of plant extract on the different bacterial strains. The MIC values showed that negative Gram bacteria (*E. coli* and *Kl. oxytoca*) were inhibited at 0.25 mg/ml and the positive Gram bacteria (*S. aureus* and *S. epidermis*) were inhibited at 0.50 mg/ml. (MIC tests pictures in Supplementary data).

Table 2. Antibacterial activity of methanolic extract of *S. securidaca* seeds

Bacterial strains	Zone of inhibition (mm) / Mean \pm SD		
	2 mg/disc	1 mg/disc	0.5 mg/disc
<i>E. coli</i>	09 \pm 1.1	07 \pm 1.0	05 \pm 1.6
<i>S. aureus</i>	13 \pm 2.1	11 \pm 1.1	09 \pm 1.0
<i>S. epidermis</i>	13 \pm 1.3	11 \pm 1.0	08 \pm 1.6
<i>Kl. oxytoca</i>	08 \pm 1.2	05 \pm 1.3	04 \pm 1.5




Fig. 2. Antimicrobial activity of methanol extract of seeds of *S. securidaca* against four bacterial strains showing zone of inhibition, concentration: 2000, 1000 and 500 μ g/ml, P= positive control (TPZ), N= negative control (water)

3.4 Antioxidant Activity

The DPPH radical scavenging activity of *S. securidaca* seeds methanol extract is shown in Fig. 3. This result has been evaluated using the DPPH radical method by reference standard (Trolox). The concentration spread of 1-100 μ g/ml. The IC₅₀ value (half maximal inhibitory concentration) was 0.057 mg/ml for *S. securidaca*.

4. DISCUSSION

Recently, drug development and phyto-medicine are the hot topics in the world in order to find new

and develop the known potential antioxidant and antibacterial. Plants are becoming more valuable in these topics because they are rich in several classes of secondary metabolites like alkaloids, polyphenols, terpenoids, steroids, glycosides and other natural products. These constituents showed an important value of herbal medicine in advances clinical research in infection diseases and improve health care [29]. The Folin-Ciocalteu's method has been used to find total phenol concentration presents in the extract. This method has been applied for finding polyphenols and other interfering compounds because of their antimicrobial and antioxidant activities. These compounds allow the extract to act as




Fig. 3. Scavenging effect of *S. securidaca* seeds extract on DPPH radical

antioxidants due to their redox properties and chemical structures [30]. The activity of bacteria against alkaloids and their derivatives such as highly aromatic planar quaternary is referred to their ability to intercalate with DNA [31, 32]. The methanolic extract of *S. securidaca* seeds uptake several natural phytochemical constituents with a percentage yield of 15.4 % (Table 4). The presence of these compounds is thought to be responsible for antimicrobial and antioxidant activity.

Table 3. MIC of *S. securidaca* seeds methanolic extract (mg/ml)

Microorganism	MIC mg/ml
<i>E. coli</i>	0.25
<i>S. aureus</i>	0.50
<i>S. epidermis</i>	0.50
<i>Kl. oxytoca</i>	0.25

Table 4. Antioxidant and total phenolic content and percentage yield

Phytochemical tests	The seeds parts of <i>S. securidaca</i> methanolic extract
Antioxidant activity (IC ₅₀)	56.74 ± 1.2 µg/ml plant extract
Total phenolic contents	62.28 ± 1.7 mg/ml plant extract
Percentage yield %	15.4% plant extract

The phytochemical constituents of *S. securidaca* seeds extract have several secondary metabolites such as flavonoids and cardiac

glycosides. Furthermore, *S. securidaca* extract has some flavonoids that act as potent cytotoxicity against HT-29 (colon carcinoma), T47D (breast ductal carcinoma) and Caco-2 (colorectal adenocarcinoma) which known as Human cancer cell line [33,34]. Meanwhile, the extract had potent antioxidant activity against DPPH with investigated IC₅₀= 56.74 ± 1.2 µg/ml plant extract. Previous report [s16] showed that Petroleum ether and chloroform fraction of *S. securidaca* seeds have antimicrobial effects on the growth of *S. aureus* and *P. aeruginosa* (etheric extract) and only *S. aureus* has inhibited by chloroform extract. Moreover, methanolic extract has no microbial effect. This is the first report on methanol extract of *S. securidaca* seeds as antioxidant and antibacterial effects. Methanolic extract showed potent antibacterial activities against *S. aureus* and *S. epidermis* as Gram positive bacteria, also *E. coli* and *Kl. Oxytoca* as Gram negative bacteria with MICs 0.5, 0.5, 0.25 and 0.25 mg /ml, respectively. The extract work as a potent antimicrobial for both type of bacteria, but the inhibition zone diameter of extract against Gram positive is ranging from 8-13 mm, while it is 4-9 mm for Gram negative bacteria and it depends on the concentration of the extract.

5. CONCLUSION

Based on the findings of this experimental work, it can be concluded that *S. securidaca* seeds extract is a potent source of antioxidant and antimicrobial agents against Gram positive and

Gram negative bacterial strains. So that, it could be used as natural antibacterial and antioxidant agent.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Al-Quran SA, Al-Mazaideh GM. Inhibitive action of chamomile extract on the corrosion of Iron: Density Functional Theory, *Mor. J. Chem.* 2018;6(1):195-202.
- Wedian F, Al-Qudah MA, Al-Mazaideh GM. Corrosion inhibition of copper by *Capparis spinosa* L. extract in strong acidic medium: Experimental and Density Functional Theory. *Int. J. Electrochem. Sci.* 2017;12: 4664-4676.
- Al-Mazaideh GM, Carbohydrates as green corrosion inhibitors of cooper: Ab initio Study, *Jord. J. Chem.* 2017;12(4):189-200.
- Al-Mazaideh GM, Ababneh TS, Abu-Shandi KH, Jamhour RMAQ, Ayaal Salman HJ, Al-Msiedeem AM, Khalil SM. DFT calculations of mesembryanthemum nodiflorum compounds as corrosion inhibitors of Aluminium. *Phy. Sci. Inter. J.* 2016;12(1):1-7.
- Al-Mazaideh GM, Al-Zereini WA, Al-Mustafa AH, Khalil SM. The effect of nitro maleimides from a marine vibrio species compounds as a source of environmentally friendly corrosion inhibitors for metals: A computational Study. *Adv. Envi. Biol.* 2016;10(8):159-168.
- Abu-Shandi K, Al-Rawashdeh B, Al-Mazaideh GM, Abu-Nameh E, Al-Amro A, Al-Soufi H, Al-Ma'abreh A, Al-Dawdeyah A, A novel strategy for the identification of the medicinal natural products in *rubus fruticosus* plant by Using GC/MS technique: A study on leaves, stems and roots of the plant. *Adv. Anal. Chem.* 2015;5(2):31-41.
- Anesini C, Perez C, Screening of plants used in argentine folkmedicine for antimicrobial activity. *J. Ethnopharmacol.* 1993;39(2):119-128.
- Gharaman A, Flore de iran en couleurs naturelle, Tehran university press, Tehran. 1993;Vol12:1478.
- Al-Hachim GM, Maki B. Effect of *Securigera securidaca* on electroshock seizure threshold in mice. *Psychological Reports.* 1969;24:551-553.
- Hosseinzadeh H, Ramezani M, Danaei AR. Antihyperglycaemic effect and acute toxicity of *Securigera securidaca* L. seed extracts in mice. *Phytotherapy Research.* 2002;16:745-747.
- Behbahani M, Sayedipour S, Pourazar A, Shanehsazzadeh M. *In vitro* anti-HIV-1 activities of kaempferol and kaempferol-7-Oglucoside isolated from *Securigera securidaca*. *Res. Pharm. Sci.,* Decenber. 2014;9(6):463-469.
- Garjani A, Fathiazad F, Zakheri A, et al. The effect of total extract of *Securigera securidaca* L. seeds on serum lipid profiles, antioxidant status, and vascular function in hypercholesterolemic rats, *J. Ethnopharmacology.* 2009;126(3): 525-532.
- Garjani A, Fathiazad F, Zakheri A, Allaf Akbari N, Azarmie Y, Fakhrajoo A et al. The effect of total extract of *Securigera securidaca* L. seeds on serum lipid profiles, antioxidant status, and vascular function in hypercholesterolemic rats. *J.Ethnopharmacol.* 2009;126:525-32.
- Zatula VV, Kovalev IP, Kolesnikov DG. Configuration of securigenin and securigenol. *Chem. Nat Compd.* 19695:111-112.
- Komissarenko AN, Kovalev VN, Hydroxycoumarins and flavones of *Securigera securidaca*, *Chem. Nat. Comp.* 1987;23(2):252-252.
- Choi CW, Kim SC, Hwang SS, Choi BK, Ahn HJ, Lee MY, Park SH, Kim SK. Antioxidant activity and free radical scavenging capacity between Korean medicinal plants and flavonoids by assay-guided comparison. *Plant Science.* 2002;163:1161-1168.
- Sadat-Ebrahimi S, Mir MH, Amin G, Hajimehdipoor H. Identification of amino acids in *Securigera securidaca*, a popular

- medicinal herb in Iranian folk medicine. Res. J. Pharmacogn. 2014;1(1):23-26.
18. Tofighi Z, Molazem M, Doostdar B, Taban P, Shahverdi A, Samadi N, Yassaa N. Antimicrobial activities of three medicinal plants and investigation of flavonoids of *Tripleurospermum disciforme*, Iranian J. Pharm. Res. 2015; 14(1):225-231.
 19. Jaradat N, Abualhasan M, Ali I. Comparison of anti-oxidant activities and exhaustive extraction yields between wild and cultivated *Cyclamen persicum*, *Malva sylvestris* and *Urtica pilulifera* leaves. J. Appl. Pharm Sci. 2015;5(4):101-107.
 20. Jaradat N, Hussen F, Al Ali A. Preliminary phytochemical screening, quantitative estimation of total flavonoids, total phenols and antioxidant activity of *Ephedra alata* decne. J. Mater Environ. Sci. 2015;6(6): 1771-1779.
 21. Evans WC. Trease and Evans' pharmacognosy. United States: Elsevier Health Sciences; 2009.
 22. Kumar S, Rajesekaran T, Shanmugam P, Kumar VM. Phytochemical constituents and antibacterial activities of *Elaeocarpus ganitrus* Roxb. and *canthium parviflora* linn. leaves. Advanced Biotech. 2007;6: 23-25.
 23. Malik CP, Singh MB. Plant enzymology and histo-enzymology: A text Manual. Kalyani Publishers, New Delhi. 1980;286.
 24. Sakanaka S, Tachibana Y, Okada Y. Preparation and antioxidant properties of extracts of Japanese persimmon leaf tea (kakinoha-cha). Food Chemistry. 2005; 89(4):569–575.
 25. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disc method. Amer. J. Clin. Pathol. 1966;45(4):493-496.
 26. Oreky NS, Nakahara K. Antibacterial activity of extracts from some edible plants commonly consumed. Asia International J. Food Microbiology. 2003;80:223-230.
 27. Dharajiya D, Patel P, Patel M, Moitra N. *In vitro* antimicrobial activity and qualitative phytochemical analysis of *Ania somnifera* (L.) dunal extracts. Int. J. Pharm Sci. Rev. Res. 2014;27:349-54.
 28. Kaviarasan S, Naik GH, Gangabhairathi R, Anuradha CV, Priyadarsini KI. *In vitro* studies on antiradical and antioxidant activities of fenugreek (*Trigonella foenum-graecum*) seeds. Food Chemistry. 2007; 103(1):31–37.
 29. Chung YC, Chang C, Chao WW, Lin CF, Chou ST. Antioxidative activity and safety of the 50 ethanolic extracts from red bean fermented by *Bacillus subtilis* MR-K1. J. Agricultural and Food Chemistry. 2002; 50(8):2454-2458.
 30. Ganguly R, Mishra P, Sharma A. Microbes and infection. Indian, J. Microbiol. 2001;41:211-213.
 31. Igbiosa OO, Igbiosa EO, Aiyegoro OA. Antimicrobial activity and phytochemical screening of stem bark extracts from *Jatropha curcas* (Linn), Afr. J. Pharmacol. 2009;3:58-62.
 32. Kumar GS, Jayaveera KN, Kumar ACK, Sanjay UP, Swamy BMV and Kumar DVK. Antimicrobial effects of acne-inducing bacteria. Tropical, J. Pharma. Res. 2007;6: 717-723.
 33. Yadav R, Tiwari R, Chowdhary P, Pradhan C. A pharmacognostical monograph of *Trigonella foenum-graecum* seeds. Int. J. Pharm. Pharm. Sci. 2011; 3(5):442-445.
 34. Komissarenko AN, Kovalev VN. Hydroxycoumarins and flavones of *Securigera securidaca*. Chem. Nat. Compd. 1987;23:252.

© 2020 Al-Mazaideh and Al-Quran; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/58374>